

The American College of Radiology, with more than 30,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice guidelines and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice guidelines and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice guideline and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review, requiring the approval of the Commission on Quality and Safety as well as the ACR Board of Chancellors, the ACR Council Steering Committee, and the ACR Council. The practice guidelines and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice guideline and technical standard by those entities not providing these services is not authorized.

Revised 2007 (Res. 26)\*

## **PRACTICE GUIDELINE FOR THE PERFORMANCE OF SONOHYSTEROGRAPHY**

### **PREAMBLE**

These guidelines are an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. They are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care. For these reasons and those set forth below, the American College of Radiology cautions against the use of these guidelines in litigation in which the clinical decisions of a practitioner are called into question.

The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the physician or medical physicist in light of all the circumstances presented. Thus, an approach that differs from the guidelines, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in the guidelines when, in the reasonable judgment of the practitioner, such course of action is indicated by the condition of the patient, limitations of available resources, or advances in knowledge or technology subsequent to publication of the guidelines. However, a practitioner who employs an approach substantially different from these guidelines is advised to document in the patient record information sufficient to explain the approach taken.

The practice of medicine involves not only the science, but also the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment.

Therefore, it should be recognized that adherence to these guidelines will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The sole purpose of these guidelines is to assist practitioners in achieving this objective.

### **I. INTRODUCTION**

The clinical aspects contained in specific sections of this guideline (Introduction, Indications and Contraindications, Specifications of the Examination, and Equipment Specifications) were developed collaboratively by the American College of Radiology (ACR), the American Institute of Ultrasound in Medicine (AIUM), and the American College of Obstetricians and Gynecologists (ACOG). Recommendations for physician qualifications, written request for the examination, procedure documentation, and quality control may vary among the three organizations and are addressed by each separately.

This guideline has been developed to assist qualified physicians performing sonohysterography. Properly performed sonohysterography can provide information about the uterus, endometrium, and fallopian tubes. Additional studies may be necessary for complete diagnosis. Adherence to the following guideline serves to maximize the diagnostic benefit of sonohysterography.

### **II. DEFINITION**

Most clinical experience and medical literature to date have focused on the sonographic imaging of the uterus,

and specifically the endometrial cavity, using the transcervical injection of sterile fluid. Thus, terms such as saline infusion sonohysterography or simply sonohysterography have been used to describe this technique. More recently, fluids other than saline have been used for intrauterine injection. Also, emerging research is being developed in assessing tubal patency using uterine infusion of fluid.

### **III. GOAL**

The goal of sonohysterography is to visualize the endometrial cavity in more detail than is possible with routine endovaginal ultrasound.

### **IV. INDICATIONS AND CONTRAINDICATIONS**

#### **A. Indications**

The most common indication for sonohysterography is abnormal uterine bleeding in both premenopausal and postmenopausal women. Other indications include but are not limited to evaluation of:

1. Infertility and habitual abortion.
2. Congenital abnormalities of the uterine cavity.
3. Assessment of uterine cavity, especially with regard to uterine myomas, polyps, and synechiae.
4. Abnormalities detected on endovaginal sonography, including focal or diffuse endometrial or intracavitary abnormalities.
5. Endometrium suboptimally imaged by endovaginal sonography.

#### **B. Contraindications**

Sonohysterography should not be performed in a woman who is pregnant or who could be pregnant. This is usually avoided by scheduling the examination in the follicular phase of the menstrual cycle, after menstrual flow has essentially ceased, but before the patient has ovulated. In a patient with regular cycles, sonohysterography should not in most cases be performed later than the tenth day of the menstrual cycle. Sonohysterography should not be performed in patients with a pelvic infection or unexplained pelvic tenderness, which could be due to pelvic inflammatory disease. Active vaginal bleeding is not a contraindication to the procedure but may make the interpretation more challenging.

### **V. QUALIFICATIONS AND RESPONSIBILITIES OF THE PHYSICIAN**

See the [ACR Practice Guideline for Performing and Interpreting Diagnostic Ultrasound Examinations](#).

In addition, the physician shall have spent a minimum of 3 months in documented formal training in the performance, interpretation, and reporting of examinations of the female reproductive system. Additionally, the physician should supervise and interpret examinations of the female reproductive system on a regular basis. The physician should be familiar with techniques of cervical cannulation.

### **VI. WRITTEN REQUEST FOR THE EXAMINATION**

The written or electronic request for sonohysterography should provide sufficient information to demonstrate the medical necessity of the examination and allow for its proper performance and interpretation.

Documentation that satisfies medical necessity includes 1) signs and symptoms and/or 2) relevant history (including known diagnoses). Additional information regarding the specific reason for the examination or a provisional diagnosis would be helpful and may at times be needed to allow for the proper performance and interpretation of the examination.

The request for the examination must be originated by a physician or other appropriately licensed health care provider. The accompanying clinical information should be provided by a physician or other appropriately licensed health care provider familiar with the patient's clinical problem or question and consistent with the state's scope of practice requirements. (ACR Resolution 35, adopted in 2006)

### **VII. SPECIFICATIONS FOR INDIVIDUAL EXAMINATIONS**

#### **A. Patient Preparation**

Pelvic organ tenderness should be assessed during the preliminary endovaginal sonogram. If adnexal tenderness or pain suspicious for active pelvic infection is found prior to fluid infusion, the examination should be deferred until after an appropriate course of treatment. In the presence of nontender hydrosalpinges, consideration may be given to administering antibiotics at the time of the examination; in this case it is prudent to discuss the antibiotic regimen with the referring physician. A pregnancy test is advised when clinically indicated. Patients should be questioned about latex allergy prior to use of a latex sheath. The optimal time to perform this test in a menstruating woman is after the bleeding ends but prior to ovulation.

#### **B. Procedure**

Preliminary routine endovaginal sonography with measurements of endometrium and evaluation of the

uterus and ovaries should be performed prior to sonohysterography. A speculum is used to allow visualization of the cervix. The presence of unusual pain, lesions, or purulent vaginal or cervical discharge may require rescheduling the procedure pending further evaluation. After cleansing the external os, the cervical canal and/or uterine cavity should be catheterized using aseptic technique, and appropriate sterile fluid should be instilled slowly by means of manual injection under real-time sonographic imaging. Imaging should include real-time scanning of the endometrial and cervical canal.

#### C. Contrast Agent

Appropriate sterile fluid such as normal saline or water should be used for sonohysterography.

#### D. Images

Appropriate images, in at least two planes, using a high-frequency endovaginal ultrasound probe should be produced and recorded to demonstrate normal and abnormal findings. Precatheterization images should be obtained, including the thickest bi-layer endometrial measurement on a sagittal image.

Once the uterine cavity is filled with fluid, representative images with a complete survey of the uterine cavity are obtained as necessary for diagnostic evaluation. If a balloon catheter is employed for the examination, images should be obtained at the end of the procedure with the balloon deflated to fully evaluate the endometrial cavity, particularly the cervical canal and lower uterine segment.

### VIII. DOCUMENTATION

Adequate documentation is essential for high quality in patient care. There should be a permanent record of the ultrasound examination and its interpretation. Comparison with prior relevant imaging studies may prove helpful. Images of all appropriate areas, both normal and abnormal, should be recorded. Variations from normal size should generally be accompanied by measurements. Images should be labeled with the patient identification, facility identification, examination date, and image orientation. An official interpretation (final report) of the ultrasound examination should be included in the patient's medical record. Retention of the ultrasound examination images should be consistent both with clinical need and with relevant legal and local health care facility requirements.

Reporting should be in accordance with the [ACR Practice Guideline for Communication of Diagnostic Imaging Findings](#).

### IX. EQUIPMENT SPECIFICATIONS

Sonohysterography is usually conducted with an endovaginal transducer. In cases of an enlarged uterus, additional transabdominal images during infusion may be required to fully evaluate the endometrium. The transducer should be adjusted to operate at the highest clinically appropriate frequency under the ALARA (as low as reasonably achievable) principle.

### X. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION

All transducers should be cleaned after use. Vaginal transducers should be covered by a protective sheath prior to insertion. Sterile coupling gel should be used. Following the examination, the sheath should be disposed of and the transducer cleaned in an antimicrobial solution. The type of solution and amount of time for cleaning depend on manufacturer and infectious disease control recommendations.

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education Concerns appearing elsewhere in the ACR Practice Guidelines and Technical Standards book.

Equipment performance monitoring should be in accordance with the [ACR Technical Standard for Diagnostic Medical Physics Performance Monitoring of Real Time Ultrasound Equipment](#).

### ACKNOWLEDGEMENTS

This guideline was revised according to the process described in the ACR Practice Guidelines and Technical Standards book by the ACR Guidelines and Standards Committee of the Commission on Ultrasound in collaboration with the American Institute of Ultrasound in Medicine (AIUM) and the American College of Obstetricians and Gynecologists (ACOG).

#### Collaborative Subcommittee

##### ACR

Gretchen A. Gooding, MD, Chair  
Ruth B. Goldstein, MD  
Mindy M. Horrow, MD

##### AIUM

Anna Lev-Toaff, MD  
Anna K. Parsons, MD  
Eugene C. Toy, MD

##### ACOG

Daniel Breitkopf, MD  
Steven R. Goldstein, MD

### ACR Guidelines and Standards Committee

Gretchen A. Gooding, MD, Chair  
Raymond E. Bertino, MD  
Mary C. Frates, MD  
Ruth B. Goldstein, MD  
Beatrice L. Madrazo, MD  
Jon W. Meilstrup, MD  
Michelle L. Melany, MD  
Miriam N. Mikhail, MD  
Sara M. O'Hara, MD  
Suhās G. Parulekar, MD  
John S. Pellerito, MD  
Philip W. Ralls, MD  
Michelle L. Robbin, MD  
Carol M. Rumack, MD, Chair, Commission

### Comments Reconciliation Committee

Jon D. Shanser, MD, Co-Chair  
Bill H. Warren, MD, Co-Chair  
Mary C. Frates, MD  
Ruth B. Goldstein, MD  
Gretchen A. Gooding, MD  
Mindy M. Horrow, MD  
Paul A. Larson, MD  
Lawrence A. Liebscher, MD  
Carol M. Rumack, MD  
Julie K. Timins, MD  
William G. Way, Jr., MD

### Suggested Reading

1. Becker E Jr, Lev-Toaff AS, Kaufman EP, Halpern EJ, Edelweiss MI, Kurtz AB. The added value of transvaginal sonohysterography over transvaginal sonography alone in women with known or suspected leiomyoma. *J Ultrasound Med* 2002;21:237-247.
2. Benacerraf BR, Shipp TD, Bromley B. Improving the efficiency of gynecologic sonography with 3-dimensional volumes: a pilot study. *J Ultrasound Med* 2006;25:165-171.
3. Bree RL, Bowerman RA, Bohm-Velez M, et al. US evaluation of the uterus in patients with postmenopausal bleeding: a positive effect on diagnostic decision making. *Radiology* 2000;216:260-264.
4. Breikopf DM, Frederickson RA, Snyder RR. Detection of benign endometrial masses by endometrial stripe measurement in premenopausal women. *Obstet Gynecol* 2004;104:120-125.
5. Doubilet PM. Society of Radiologists in Ultrasound Consensus Conference statement on postmenopausal bleeding. *J Ultrasound Med* 2001;20:1037-1042.
6. Dubinsky TJ, Stroehlein K, Abu-Ghazze Y, Parvey HR, Maklad N. Prediction of benign and malignant endometrial disease: hysterosonographic-pathologic correlation. *Radiology* 1999;210:393-397.
7. Goldstein RB, Bree RL, Benson CB, et al. Evaluation of women with postmenopausal bleeding: Society of

Radiologists in Ultrasound-Sponsored Consensus Conference statement. *J Ultrasound Med* 2001;20:1025-1036.

8. Goldstein SR. Use of ultrasonohysterography for triage of perimenopausal patients with unexplained uterine bleeding. *Am J Obstet Gynecol* 1994;170:565-570.
9. Hann LE, Gretz EM, Bach AM, Francis SM. Sonohysterography for evaluation of the endometrium in women treated with tamoxifen. *AJR* 2001;177:337-342.
10. Laifer-Narin SL, Ragavendra N, Lu DS, Sayre J, Perrella RR, Grant EG. Transvaginal saline hysterosonography: characteristics distinguishing malignant and various benign conditions. *AJR* 1999;172:1513-1520.
11. Laifer-Narin SL, Ragavendra N, Parmenter EK, Grant EG. False-normal appearance of the endometrium on conventional transvaginal sonography: comparison with saline hysterosonography. *AJR* 2002;178:129-133.
12. Lev-Toaff AS, Toaff ME, Liu JB, Merton DA, Goldberg BB. Value of sonohysterography in the diagnosis and management of abnormal uterine bleeding. *Radiology* 1996;201:179-184.
13. Lindheim SR, Sprague C, Winter TC 3rd. Hysterosalpingography and sonohysterography: lessons in technique. *AJR* 2006;186:24-29.
14. Mihm LM, Quick VA, Brumfield JA, Connors AF Jr, Finnerty JJ. The accuracy of endometrial biopsy and saline sonohysterography in the determination of the cause of abnormal uterine bleeding. *Am J Obstet Gynecol* 2002;186:858-860.
15. Parsons AK, Lense JJ. Sonohysterography for endometrial abnormalities: preliminary results. *J Clin Ultrasound* 1993;21:87-95.
16. Schwartz LB, Snyder J, Horan C, Porges RF, Nachtigall LE, Goldstein SR. The use of transvaginal ultrasound and saline infusion sonohysterography for the evaluation of asymptomatic postmenopausal breast cancer patients on tamoxifen. *Ultrasound Obstet Gynecol* 1998;11:48-53.

---

\*Guidelines and standards are published annually with an effective date of October 1 in the year in which amended, revised or approved by the ACR Council. For guidelines and standards published before 1999, the effective date was January 1 following the year in which the guideline or standard was amended, revised, or approved by the ACR Council.

### Development Chronology for this Guideline

2002 (Resolution 28)  
Amended 2006 (Resolution 35)  
Revised 2007 (Resolution 26)